

Modern Physics Letters A
 © World Scientific Publishing Company

EXTENSION OF GRIMUS-STOCKINGER FORMULA FROM OPERATOR EXPANSION OF FREE GREEN FUNCTION

S. E. Korenblit and D. V. Taychenachev
*Department of Physics, Irkutsk State University,
 Gagarin blvd 20, Irkutsk 664003, Russia
 korenblit@ic.isu.ru*

Received (Day Month Year)
 Revised (Day Month Year)

The operator expansion of free Green function of Helmholtz equation for arbitrary N - dimension space leads to asymptotic extension of 3- dimension Grimus-Stockinger formula closely related to multipole expansion. Analytical examples inspired by neutrino oscillation and neutrino deficit problems are considered for relevant class of wave packets.

Keywords: asymptotic expansion; Green function; wave packet; neutrino deficit.

PACS Nos.: 03.65.Db, 11.80.Fv, 14.60.Pq

1. Introduction

A considerable efforts was made recently¹⁻³ to extend so called Grimus-Stockinger theorem,⁴ which is the main tool of the modern theory of neutrino oscillations⁵⁻⁷ and gives the leading asymptotic behavior with $R = |\mathbf{R}| \rightarrow \infty$ for the integral:

$$\mathcal{J}(\mathbf{R}) = \int \frac{d^3q}{(2\pi)^3} \frac{e^{-i(\mathbf{q}\cdot\mathbf{R})} \Phi(\mathbf{q})}{(\mathbf{q}^2 - k^2 - i0)} \sim \frac{e^{ikR}}{4\pi R} \Phi(-k\mathbf{n}) \left[1 + O(R^{-1/2}) \right], \quad (1)$$

where: $\mathbf{R} = R\mathbf{n}$, $\mathbf{n}^2 = 1$, and the function $\Phi(\mathbf{q}) \in C^3$ decreases at least like $1/\mathbf{q}^2$ together with its first and second derivatives. In the next section of the present letter it is reminded that the possibility of further asymptotic expansion and the order of leading correction depend⁸ on the chosen properties^{1,2} of this function. In section 3 for appropriate space of $\Phi(\mathbf{q})$ a closed formula and simple recurrent relation for coefficients of asymptotic expansion of $\mathcal{J}(\mathbf{R})$ in all orders of R^{-s} are obtained. Such extension has particular importance for explanation^{2,3} of observed⁹ (anti-) neutrino deficit at short distances from the sources discussed in section 4. But it may find much more wide implementation in quantum physics and optics, when the Green function of Helmholtz equation is used. In sections 4, 5 the space chosen for the functions $\Phi(\mathbf{q})$ is advocated on the ground of quantum field theory of wave packets.^{6,10} Some useful relations and generalization onto N - dimension case are placed in Appendix.

2. Preliminaries

In order to understand the physical nature of asymptotic expansion, we notice that for infinitely differentiable $\Phi(\mathbf{q})$ uniquely representable by its Taylor expansion for any finite $|\mathbf{q}| < \infty$: $e^{-i(\mathbf{q} \cdot \mathbf{R})} \Phi(\mathbf{q}) = \Phi(i \nabla_{\mathbf{R}}) e^{-i(\mathbf{q} \cdot \mathbf{R})}$, and then, formally:

$$\mathcal{J}(\mathbf{R}) = \Phi(i \nabla_{\mathbf{R}}) \frac{e^{ikR}}{4\pi R} = \Phi(-i \nabla_{\mathbf{x}}) \frac{e^{ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} \Big|_{\mathbf{x}=0}, \quad (2)$$

where the differential vector operator in spherical basis $\mathbf{n}, \boldsymbol{\eta}_\vartheta, \boldsymbol{\eta}_\varphi$ has the following properties:

$$\mathbf{n} = (\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta), \quad \boldsymbol{\eta}_\vartheta = \partial_\vartheta \mathbf{n}, \quad \sin \vartheta \boldsymbol{\eta}_\varphi = \partial_\varphi \mathbf{n}, \quad (3)$$

$$\nabla_{\mathbf{R}} = \mathbf{n} \partial_R + \frac{1}{R} \partial_{\mathbf{n}}, \quad (\mathbf{n} \cdot \nabla_{\mathbf{R}}) = \partial_R, \quad \partial_{\mathbf{n}} = \boldsymbol{\eta}_\vartheta \partial_\vartheta + \frac{\boldsymbol{\eta}_\varphi}{\sin \vartheta} \partial_\varphi, \quad (4)$$

$$(\mathbf{n} \cdot \partial_{\mathbf{n}}) = 0, \quad (\partial_{\mathbf{n}} \cdot \mathbf{n}) = 2, \quad (\mathbf{n} \times \partial_{\mathbf{n}})^2 = \partial_{\mathbf{n}}^2, \quad (\mathbf{n} \times \partial_{\mathbf{n}}) = i \mathbf{L}_{\mathbf{n}}, \quad (5)$$

$$-\partial_{\mathbf{n}}^2 = \mathbf{L}_{\mathbf{n}}^2 = 2R(\mathbf{n} \cdot \nabla_{\mathbf{R}}) + R^2((\mathbf{n} \cdot \nabla_{\mathbf{R}})^2 - \nabla_{\mathbf{R}}^2), \quad \text{whence,} \quad (6)$$

$$\text{for } \cos \vartheta = c: \quad \mathbf{L}_{\mathbf{n}}^2 = -[\partial_c(1-c^2)\partial_c + (1-c^2)^{-1}\partial_\varphi^2] \equiv \mathcal{L}_{\mathbf{n}}, \quad (7)$$

and the well known representation at point \mathbf{R} for the spherical wave coming from point \mathbf{x} , as a free Schrödinger's 3-dimensional Green function,¹¹ is used (see (A.6)):

$$\frac{e^{ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} = \int \frac{d^3\mathbf{q}}{(2\pi)^3} \frac{e^{\pm i(\mathbf{q} \cdot (\mathbf{R}-\mathbf{x}))}}{(\mathbf{q}^2 - k^2 - i0)}, \quad (8)$$

which for $\mathbf{x} = 0$ satisfies to the equation:

$$(-\nabla_{\mathbf{R}}^2 - k^2) \frac{e^{\pm ikR}}{4\pi R} = \delta_3(\mathbf{R}), \quad (9)$$

where for $R > 0$ $\nabla_{\mathbf{R}}^2$ is given also by (A.1) with $N = 3$. Since for $R > 0$ the right hand side of this equation is zero, $\delta_3(\mathbf{R}) = 0$, for spherically symmetric case one immediately obtains from (2) formally an exact answer, which really takes place at least for the function $\Psi(k^2)$ regular and bounded in upper half plane $\text{Im } k \geq 0$ of complex variable k :

$$\text{when } \Phi(\mathbf{q}) = \Psi(q^2), \text{ for } \mathbf{q}^2 = q^2, \text{ then: } \mathcal{J}(\mathbf{R}) = \Psi(k^2) e^{ikR} / (4\pi R). \quad (10)$$

Therefore the higher order corrections originate only by asymmetry of function $\Phi(\mathbf{q})$ relative to the directions of \mathbf{q} in accordance with our previous result.¹ Indeed, in order to obtain them, we supposed¹ that $\Phi(\mathbf{q})$ and its first and second derivatives are represented by Fourier-transforms as:

$$\Phi(\mathbf{q}) = \int d^3\mathbf{x} e^{i(\mathbf{q} \cdot \mathbf{x})} \phi(\mathbf{x}), \quad \nabla_{\mathbf{q}} \Phi(\mathbf{q}) = i \int d^3\mathbf{x} e^{i(\mathbf{q} \cdot \mathbf{x})} \mathbf{x} \phi(\mathbf{x}), \text{ and so on.} \quad (11)$$

Then by interchanging the order of integration and using the Eq. (8) we found the representation:

$$\mathcal{J}(\mathbf{R}) = \int d^3\mathbf{x} \frac{e^{ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} \phi(\mathbf{x}). \quad (12)$$

Substituting here the expansion, which in the exponential should always contain one additional order in compare with the number of orders in denominator, for:

$$|\mathbf{R} - \mathbf{x}| = R \left[1 - 2 \frac{(\mathbf{n} \cdot \mathbf{x})}{R} + \frac{\mathbf{x}^2}{R^2} \right]^{1/2}, \quad \text{whih: } \mathbf{x}^2 - (\mathbf{n} \cdot \mathbf{x})^2 \Rightarrow (\mathbf{n} \times \mathbf{x})^2, \quad (13)$$

$$|\mathbf{R} - \mathbf{x}| = R - (\mathbf{n} \cdot \mathbf{x}) + \frac{\mathbf{x}^2 - (\mathbf{n} \cdot \mathbf{x})^2}{2R} + \dots, \quad (14)$$

we had the corresponding expansion of integral (12) up to $O(R^{-3})$:

$$\mathcal{J}(\mathbf{R}) = \frac{e^{ikR}}{4\pi R} \int d^3x e^{-ik(\mathbf{n} \cdot \mathbf{x})} \phi(\mathbf{x}) \left[1 + \frac{(\mathbf{n} \cdot \mathbf{x})}{R} + \frac{ik}{2R} (\mathbf{x}^2 - (\mathbf{n} \cdot \mathbf{x})^2) + \dots \right],$$

that by making use of (11) was immediately transcribed as:

$$\mathcal{J}(\mathbf{R}) = \frac{e^{ikR}}{4\pi R} \left[1 - \frac{i}{R} (\mathbf{n} \cdot \nabla_q) + \frac{ik}{2R} ((\mathbf{n} \cdot \nabla_q)^2 - \nabla_q^2) + \dots \right] \Phi(\mathbf{q}) \Big|_{\mathbf{q}=-k\mathbf{n}}, \quad (15)$$

$$\text{with: } (\mathbf{n} \cdot \nabla_q) \Phi(\mathbf{q}) \Big|_{\mathbf{q}=-k\mathbf{n}} = -(\mathbf{n} \cdot \nabla_k) \Phi(-\mathbf{k}) = -\partial_k \Phi(-k\mathbf{n}), \text{ and so on.} \quad (16)$$

The next observation is that due to (16), with $\mathbf{k} = k\mathbf{n}$, for $\mathbf{q} = -\mathbf{k}$, $\nabla_q = -\nabla_k$, from (6), (7) with $\mathbf{R} \mapsto \mathbf{k}$, one has: $2k(\mathbf{n} \cdot \nabla_k) + k^2 ((\mathbf{n} \cdot \nabla_k)^2 - \nabla_k^2) = \mathcal{L}_n$, and expression (15) takes the following simple form:

$$\mathcal{J}(\mathbf{R}) = \frac{e^{ikR}}{4\pi R} \left[1 + \frac{i}{R} (\mathbf{n} \cdot \nabla_k) + \frac{ik}{2R} ((\mathbf{n} \cdot \nabla_k)^2 - \nabla_k^2) + \dots \right] \Phi(-\mathbf{k}) = \quad (17)$$

$$= \frac{e^{ikR}}{4\pi R} \left[1 + \frac{i}{2kR} \mathcal{L}_n + O(R^{-2}) \right] \Phi(-k\mathbf{n}). \quad (18)$$

In the next section by using the another surprisingly simple way, this result is generalized onto all orders of R^{-s} . It should be noted, that in spite of the one and the same final results of the both Eqs. (15) and (17), (18), the substitution like (13): $\nabla_q^2 - (\mathbf{n} \cdot \nabla_q)^2 \Rightarrow (\mathbf{n} \times \nabla_q)^2$ is correct till $\mathbf{q} \neq -k\mathbf{n}$ for expressions¹⁻³ like (15), but it becomes incorrect for non-commutative operators \mathbf{n} and ∇_k in expressions like (17). This difference from Ref. 1-3 below eventually gives rise an explicit operator expression for higher coefficients of this asymptotic expansion with arbitrary s .

3. Asymptotic and multipole expansions

Lemma 1. For $\mathbf{R} = R\mathbf{n}$, $\mathbf{x} = r\mathbf{v}$, $\mathbf{v} = (\sin \beta \cos \alpha, \sin \beta \sin \alpha, \cos \beta)$, $|\mathbf{x}| = r < R$, with operator $\mathcal{L}_n = L_n^2$ (or $\mathbf{n} \mapsto \mathbf{v}$) defined by Eqs. (5)–(7) and positively defined operator $\mathcal{L}_n + \frac{1}{4} = (\Lambda_n + \frac{1}{2})^2$, so that $\Lambda_n + \frac{1}{2} = \sqrt{\mathcal{L}_n + \frac{1}{4}}$ is positively defined:

$$\frac{e^{ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} = \frac{\chi_{\Lambda_n}(-ikR)}{4\pi R} e^{-ik(\mathbf{n} \cdot \mathbf{x})} \sim \quad (19)$$

$$\sim \frac{e^{ikR}}{4\pi R} \left\{ 1 + \sum_{s=1}^{\infty} \frac{\prod_{\mu=1}^s [\mathcal{L}_n - \mu(\mu-1)]}{s!(-2ikR)^s} \right\} e^{-ik(\mathbf{n} \cdot \mathbf{x})}. \quad (20)$$

Proof. The expression (19) is a formal operator rewrite for $R > r$ of the usual multipole expansion of the Green function¹¹ (8) via the corresponding expansion of the plane wave,¹¹ given also by formulas (8.533), (8.534) of Ref. 12, with the formally introduced instead l , but not really appeared operator $l \mapsto \Lambda_{\mathbf{n}}$:

$$\frac{e^{\pm ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} = \frac{1}{kRr} \sum_{l=0}^{\infty} i^{\mp l} \chi_l(\mp ikR) \psi_{l0}(kr) \sum_{m=-l}^l Y_l^m(\mathbf{n}) Y_l^{*m}(\mathbf{v}), \quad (21)$$

$$e^{\mp i(\mathbf{k} \cdot \mathbf{x})} = \frac{4\pi}{kr} \sum_{l=0}^{\infty} i^{\mp l} \psi_{l0}(kr) \sum_{m=-l}^l Y_l^m(\mathbf{n}) Y_l^{*m}(\mathbf{v}). \quad (22)$$

Here the spherical functions $Y_l^m(\mathbf{n}) = \langle \mathbf{n} | l m \rangle$ and Legendre polynomials $P_l(\xi)$ for $\xi = c = \cos \vartheta$ or $\xi = (\mathbf{n} \cdot \mathbf{v})$, as eigenfunctions of self-adjoint operator (6), (7) on the unit sphere, satisfy to well known orthogonality, parity and completeness conditions¹¹⁻¹⁴ with delta-function $\delta_{\Omega}(\mathbf{n}, \mathbf{v})$ on the unit sphere:

$$\mathbf{L}_{\mathbf{n}}^2 Y_l^m(\mathbf{n}) = l(l+1) Y_l^m(\mathbf{n}), \quad \mathbf{L}_{\mathbf{n}}^2 P_l(\xi) = l(l+1) P_l(\xi), \quad (23)$$

$$\int d\Omega(\mathbf{n}) Y_l^m(\mathbf{n}) Y_j^{m'}(\mathbf{n}) = \delta_{lj} \delta_{mm'}, \quad \sum_{l=0}^{\infty} \frac{(2l+1)}{4\pi} P_l((\mathbf{n} \cdot \mathbf{v})) = \delta_{\Omega}(\mathbf{n}, \mathbf{v}), \quad (24)$$

$$\sum_{m=-l}^l Y_l^m(\mathbf{n}) Y_l^{*m}(\mathbf{v}) \equiv \frac{(2l+1)}{4\pi} P_l((\mathbf{n} \cdot \mathbf{v})), \quad (-1)^l Y_l^m(\mathbf{n}) = Y_l^m(-\mathbf{n}). \quad (25)$$

The solutions $\chi_l(\mp ikr)$, $\psi_{l0}(kr)$ of free radial Schrödinger equation:

$$\left[r^2 \left(\frac{1}{r} \partial_r^2 r + k^2 \right) \right] \frac{\psi_{l0}(kr)}{r} = l(l+1) \frac{\psi_{l0}(kr)}{r}, \quad (26)$$

are defined by Macdonald $K_{\lambda}(z)$ and Bessel $J_{\lambda}(y)$ functions,^{8,11-14} that for integer l i.e. half integer $\lambda = l + \frac{1}{2}$ are reduced to elementary functions:^{8,12}

$$\chi_l(bR) \equiv \left(\frac{2bR}{\pi} \right)^{1/2} K_{l+\frac{1}{2}}(bR), \quad \chi_l(bR) \Big|_{l=\text{int}} = e^{-bR} \sum_{s=0}^l \frac{(l+s)!}{s!(l-s)!(2bR)^s}, \quad (27)$$

$$\psi_{l0}(kr) \equiv \left(\frac{\pi kr}{2} \right)^{1/2} J_{l+\frac{1}{2}}(kr) \equiv \frac{1}{2i} [i^{-l} \chi_l(-ikr) - i^l \chi_l(ikr)]. \quad (28)$$

The function $K_{\lambda}(z)$ (A.3) is a whole function^{8,12} of λ^2 , that is the reason, why the above introduced in (19) well defined operator $\Lambda_{\mathbf{n}}$ does not appear explicitly.

The expression (20) is the known asymptotic series of expression (19) at $R \rightarrow \infty$, as an infinite asymptotic version^{8,12} of the sum (27) for arbitrary non-integer l , $|\arg(bR)| < 3\pi/2$. It directly results also from the substitution of finite sums (25), (27) by interchanging the order of summations, converted the Eq. (21) into the sum:

$$\frac{e^{ikR}}{4\pi R} \sum_{s=0}^{\infty} \frac{1}{s!(-2ikR)^s} \frac{1}{kr} \sum_{l=s}^{\infty} i^{-l} \frac{(l+s)!}{(l-s)!} \psi_{l0}(kr) (2l+1) P_l((\mathbf{n} \cdot \mathbf{v})), \quad (29)$$

$$\text{where : } \frac{(l+s)!}{(l-s)!} = \prod_{\mu=1}^s (l-\mu+1)(l+\mu) = \prod_{\mu=1}^s [l(l+1) - \mu(\mu-1)], \quad (30)$$

equals to zero for all missing summands with $0 \leq l \leq s-1$ automatically and due to Eqs. (23) may be factored out from the sum over l as operator product in the right hand side of Eq. (20). The formal addition of all such missing in fact zero summands with $0 \leq l \leq s-1$ completes then sum over l into the plane wave (22) and converts the expression (29) into the expansion (20).

Remark. The operator $\mathcal{L}_{\mathbf{n}}$ in Eq. (20) with the same success may be replaced by operator in square brackets of the left hand side of Eq. (26) or by the same with interchanging $r \rightleftharpoons k$. \square

Theorem 1. *Let $\Phi(\mathbf{q}) \in S(\mathbf{R}_{\mathbf{q}}^3)$, the space of functions infinitely differentiable $\forall \mathbf{q} \in \mathbf{R}_{\mathbf{q}}^3$, decrease faster than any power of $1/|\mathbf{q}|$ together with all its derivatives. Then integral $\mathcal{J}(\mathbf{R})$ (1), (12) at $R \rightarrow \infty$ admits asymptotic expansion, which has asymptotic sense⁸ even though $\phi(\mathbf{x})$ in Eq. (11) has a finite support:*

$$\mathcal{J}(\mathbf{R}) \sim \frac{e^{ikR}}{4\pi R} \Phi(-k\mathbf{n}) \left\{ 1 + \sum_{s=1}^{\infty} \frac{C_s(k, \mathbf{n})}{(-2ikR)^s} \right\}, \quad \text{with:} \quad (31)$$

$$\Phi(-k\mathbf{n})C_s(k, \mathbf{n}) = \frac{1}{s!} \prod_{\mu=1}^s [\mathcal{L}_{\mathbf{n}} - \mu(\mu-1)] \Phi(-k\mathbf{n}), \quad \text{or:} \quad (32)$$

$$\Phi(-k\mathbf{n})C_s(k, \mathbf{n}) = \frac{\mathcal{L}_{\mathbf{n}} - s(s-1)}{s} \Phi(-k\mathbf{n})C_{s-1}(k, \mathbf{n}), \quad C_0(k, \mathbf{n}) = 1, \quad (33)$$

and which is equivalent to infinite reordering of its multipole expansion:

$$\mathcal{J}(\mathbf{R}) \sim \frac{1}{4\pi R} \sum_{j=0}^{\infty} \chi_j(-ikR) \sum_{m=-j}^j B_j^m(k) Y_j^m(\mathbf{n}), \quad (34)$$

$$\text{with: } \Phi(-k\mathbf{n})C_s(k, \mathbf{n}) = \frac{1}{s!} \sum_{j=s}^{\infty} \frac{(j+s)!}{(j-s)!} \sum_{m=-j}^j B_j^m(k) Y_j^m(\mathbf{n}), \quad (35)$$

$$\text{for: } \Phi(-k\mathbf{n}) = \sum_{j=0}^{\infty} \sum_{m=-j}^j B_j^m(k) Y_j^m(\mathbf{n}), \quad \phi(\mathbf{x}) = \int \frac{d^3\mathbf{q}}{(2\pi)^3} \Phi(\mathbf{q}) e^{-i(\mathbf{q} \cdot \mathbf{x})}. \quad (36)$$

Proof. Since Fourier transformation (11) maps the space $S(\mathbf{R}^3)$ into itself¹⁵ the function $\phi(\mathbf{x}) \in S(\mathbf{R}_{\mathbf{x}}^3)$ also and is represented by the inverse Fourier transform (36). Let suppose at first that $\phi(\mathbf{x})$ has a finite support at $|\mathbf{x}| \leq r_0$. Then for $R > r_0$ we can directly substitute the expressions (20) into representation (12) of $\mathcal{J}(\mathbf{R})$ with the following result after interchange of the order of summation, differentiation and integration of Fourier transform (11), justified⁸ also for the asymptotic series:

$$\mathcal{J}(\mathbf{R}) = \frac{\chi_{\Lambda_{\mathbf{n}}}(-ikR)}{4\pi R} \Phi(-k\mathbf{n}) \sim \frac{e^{ikR}}{4\pi R} \left\{ 1 + \sum_{s=1}^{\infty} \frac{\prod_{\mu=1}^s [\mathcal{L}_{\mathbf{n}} - \mu(\mu-1)]}{s!(-2ikR)^s} \right\} \Phi(-k\mathbf{n}). \quad (37)$$

6 *S. E. Korenblit and D. V. Taychenachev*

This is exactly asymptotic expansion (31) with dimensionless coefficients $C_s(k, \mathbf{n})$, defined by Eq. (32). However that is not the case for the function $\phi(\mathbf{x})$ with infinite support. Estimating it for $r > R$ as $|\phi(\mathbf{x})| < C_M/r^M$ with arbitrary finite $M \gg 1$, the two pieces of correction that should be added, are easy estimating as:

$$\Delta_R \mathcal{J} = \int_{r>R} d^3\mathbf{x} \frac{e^{ik|\mathbf{R}-\mathbf{x}|}}{4\pi|\mathbf{R}-\mathbf{x}|} \phi(\mathbf{x}), \quad \Delta_R \Phi = -\frac{\chi_{\Lambda_n}(-ikR)}{4\pi R} \int_{r>R} d^3\mathbf{x} e^{-ik(\mathbf{n}\cdot\mathbf{x})} \phi(\mathbf{x}), \quad (38)$$

$$|\Delta_R \mathcal{J}| < \frac{C_M}{(M-2)R^{M-2}}, \quad |\Delta_R \Phi| < \frac{C_M}{(M-3)R^{M-2}} [1 + O(R^{-1})]. \quad (39)$$

Due to this corrections and arbitrariness of $M \gg 1$, the expansion (31), (37) acquires an additional asymptotic meaning⁸ in comparison with expansion (20). The terms with $s = 0, 1$ reproduce the results¹ (10), (18) of previous section.

Coefficients of multipole expansion in (36) may be defined by the following two different ways, whose equivalence, for $\mathbf{x} = r\mathbf{v}$:

$$B_l^m(k) \equiv \int d\Omega(\mathbf{n}) \Phi(-k\mathbf{n}) \dot{Y}_l^m(\mathbf{n}) = 4\pi \int d^3\mathbf{x} \frac{\psi_{l0}(kr)}{kr} i^{-l} \dot{Y}_l^m(\mathbf{v}) \phi(\mathbf{x}), \quad (40)$$

follows from Eqs. (11), (22)–(25) and (A.4), with inverse Fourier transform (36).

Substitution of multipole expansion of Green function (21) into the integral (12) by making use of definition (40), after the same steps and under the same conditions for $\phi(\mathbf{x})$ as above leads to the **multipole** expansion (34), which with the help of expansion (27) transcribes again as asymptotic expansion (31) with the coefficients given by Eq. (35). But the same expression (35) is obtained by direct substitution of (36) into the definition (32) by means of (23), (30). This confirms the equivalence of expansions (31) and (34).

Corollary 1. From (31) with $C_s = C_s(k, \mathbf{n})$, for the squared absolute value follows:

$$(4\pi R)^2 |\mathcal{J}(\mathbf{R})|^2 \sim \sim |\Phi(-k\mathbf{n})|^2 \left\{ 1 + \sum_{s=1}^{\infty} \frac{i^s [C_s + (-1)^s C_s^*]}{(2kR)^s} + \sum_{\zeta=2}^{\infty} \frac{i^{\zeta}}{(2kR)^{\zeta}} \sum_{s=1}^{\zeta-1} (-1)^{\zeta-s} C_s^* C_{\zeta-s} \right\}. \quad (41)$$

For $\Phi(-\mathbf{k}) = \Phi^*(-\mathbf{k})$: $C_s = C_s^*$, so $s \mapsto 2n$ at the first sum over s . But the second internal sum over $1 \leq s \leq \zeta - 1$ in (41) is equal to itself with the multiplier $(-1)^{\zeta}$ and thus $\zeta \mapsto 2n$ also. Dividing further this sum over $1 \leq s \leq 2n - 1$ into two parts: with $1 \leq s \leq n$ and $n + 1 \leq s \leq 2n - 1$, and putting for the second sum $s = 2n - s'$, one finds:

$$(4\pi R)^2 |\mathcal{J}(\mathbf{R})|^2 \sim \Phi^2(-k\mathbf{n}) \left\{ 1 + \sum_{n=1}^{\infty} \frac{\Upsilon_n}{(2kR)^{2n}} \right\}, \quad \text{with: } \Upsilon_n = \Upsilon_n(k, \mathbf{n}), \quad (42)$$

$$\Upsilon_n = (-1)^n \left[2C_{2n} + \sum_{s=1}^{2n-1} (-1)^s C_s C_{2n-s} \right] = (-1)^n \sum_{s=0}^{2n} (-1)^s C_s C_{2n-s}, \quad (43)$$

$$\text{or: } \Upsilon_n = 2(-1)^n \sum_{s=0}^n (-1)^s C_s C_{2n-s} - (C_n)^2, \quad \Upsilon_1 = (C_1)^2 - 2C_2, \quad (44)$$

what, due to “homogeneity” over C_s relative to index s “like a power”, coincides with the relations² for dimensional analogs of coefficients C_s and Υ_s .

Corollary 2. Asymptotic expansion of the integral $\tilde{\mathcal{J}}(\mathbf{R}) = \mathcal{J}(-\mathbf{R})$ (1) is obtained by substitutions $\mathbf{n} \mapsto -\mathbf{n}$, $\mathcal{L}_{-\mathbf{n}} = \mathcal{L}_{\mathbf{n}}$ (7) in Eqs. (31)–(33), (37), (41)–(44). \square

The explicit Eqs. (31), (32) and recurrent relation (33) may be compared with seminumerical calculations^{2,3} of the same expansion, that for arbitrary $\Phi(\mathbf{q}) \in S(\mathbf{R}_{\mathbf{q}}^3)$ have to give the same coefficients for the same orders R^{-s} . For the first correction this is easily seen from Eqs. (15)–(18) and the Eq. (8) of Ref. 2 respectively with substitution $\nabla_q^2 - (\mathbf{n} \cdot \nabla_q)^2 \rightleftharpoons (\mathbf{n} \times \nabla_q)^2$. The same takes place for the next few corrections computed there. It is clear that any common multiplicative dependence of function $\Phi(\mathbf{q})$ on $|\mathbf{q}| = k$ does not affect the coefficients $C_s(k, \mathbf{n})$ (32) and $\Upsilon_n(k, \mathbf{n})$ (43). But there are no ways to trace this property for arbitrary order s with the only numerically calculable coefficients in Eqs. (41)–(46) of Ref. 2. An important advantage of the results (31)–(33) for coefficients of asymptotic expansion is not only the explicit arbitrariness of their order s , but the elucidating, that in fact they are defined by dependence of $\Phi(\mp k\mathbf{n})$ on the *unit vector \mathbf{n} only*. This essentially reduces the number of possible degrees of freedom and simplifies the further analytical consideration. Indeed, when $B_l^m(k) = 0$ for $l > j$, the dependence (33), (35), (44) of Υ_s on function $\Phi(\mathbf{q})$ (36) leads to termination of the asymptotic expansion: $C_s = 0 = \Upsilon_s$, for $s > j$, which is also clear from Eq. (34). If one has:

$$\Phi(-k\mathbf{n}) = \sum_{m=-j}^j B_j^m(k) Y_j^m(\mathbf{n}), \text{ then: } C_n = \frac{(j+n)!}{n!(j-n)!}, \quad \Upsilon_1 = 2j(j+1) > 0, \quad (45)$$

$$\Upsilon_j = (C_j)^2, \quad \Upsilon_n = \frac{2(-1)^n}{(2n)!} \sum_{s=0}^n (-1)^s C_{2n}^s \frac{(j+s)!(j+2n-s)!}{(j-s)!(j-2n+s)!} - (C_n)^2, \quad (46)$$

where $0 \leq n \leq j$, $\max(0, 2n-j) \leq s \leq n$ and C_{2n}^s are binomial coefficients.

4. Neutrino deficit and wave packets

At diagrammatic treatment of neutrino oscillation²⁻⁷ the function $\Phi(\mathbf{q})$ in Eq. (1) represents an overlap function $\mathcal{F}(q) = \mathcal{F}_C(q)\mathcal{F}_D(q)$ defined for $q^\mu = (q^0, \mathbf{q})$ as a real-valued product of convolutions of the wave packets of external particles participating in the processes of (anti-) neutrino creation $\{C\}$ and detection $\{D\}$ at respective space-time points. According to the Feynman rules for respective tree amplitude⁴⁻⁷ one has to deal with asymptotic expansion of $\mathcal{J}(\mathbf{R})$ (1) for the case of antineutrino⁴ or with expansion of $\tilde{\mathcal{J}}(\mathbf{R})$ for the case of neutrino,⁵ where the vector $\mathbf{R} = R\mathbf{n} = \mathbf{X}_D - \mathbf{X}_C$ for both cases defines the macroscopic geometrical parameters of observation:⁵ distance R and direction \mathbf{n} . The event rate is obtained to be proportional to $|\mathcal{J}(\mathbf{R})|^2$ or $|\tilde{\mathcal{J}}(\mathbf{R})|^2$ respectively (see for detail Refs. 2,5,6,7) that due to (42) in the leading and next to leading orders reads:

$$\propto \frac{\Phi^2(\mp k\mathbf{n})}{R^2} \left\{ 1 + \frac{\Upsilon_1(k, \pm\mathbf{n})}{(2kR)^2} + \dots \right\}. \quad (47)$$

Here the absolute value of momentum $k = \sqrt{q_0^2 - m_j^2}$ is defined by neutrino mass m_j and its mean energy $q^0 = \mp Q^0$ near its most probable energies $Q_C^0 \approx Q_D^0$, defined in turn⁴⁻⁷ by fixed parameters of respective external wave packets, such as their masses m_a , 4-vectors of most probable energy-momentum p_a and corresponding widths $\sigma_a \ll m_a$, because in the plane-wave limit, $\sigma_a \rightarrow 0$, $\forall a$ the overlap function keeps exact energy-momentum conservation for both vertices, fully destroying the neutrino-oscillations pattern,⁵ $\mathcal{F}(q)|_{\sigma_a \rightarrow 0} \propto \delta_4(q \pm Q_C)\delta_4(q \pm Q_D)$, where Q_C becomes then a full 4-momentum incoming into (anti-) neutrino propagator, whereas Q_D is full its outgoing 4-momentum. For $\sigma_a > 0$ the function $\mathcal{F}(q)$ can provide thus only approximate or “smeared” conservation⁶ of some mean (anti-) neutrino energy $Q^0 \approx Q_C^0 \approx Q_D^0$ and some mean (anti-) neutrino 3-momentum $\mathbf{Q} \approx \mathbf{Q}_C \approx \mathbf{Q}_D$, whose explicit form depends on the chosen models of wave packets and the used approximations.⁵⁻⁷ So, the function $\Phi(\mp k\mathbf{n}) = \mathcal{F}(q^0, \mathbf{q})|_{\mathbf{q}=\mp k\mathbf{n}}$ may be more or less sharply peaked^{5,6} and varies rapidly: near the point $k\mathbf{n} \simeq \mathbf{Q} = |\mathbf{Q}|\boldsymbol{\rho}$, for the more rough case⁵ (a); or near the two adjacent points $k\mathbf{n} \simeq \mathbf{Q}_{C,D} = |\mathbf{Q}_{C,D}|\boldsymbol{\rho}_{C,D}$, where $\boldsymbol{\rho}_{C,D}^2 = 1$, for the more precisely case⁶ (b). For the case (a), up to fully unessential now multiplicative dependence on k we remain with a function $\Phi(\mp k\mathbf{n}) \mapsto f(\xi)$ of the one dimensionless variable $\xi = (\boldsymbol{\rho} \cdot \mathbf{n})$ only. For the case (b) without full symmetry relative to vectors $\boldsymbol{\rho}_{C,D}$ we remain with a function $\Phi(\mp k\mathbf{n}) \mapsto \mathcal{W}(\xi_C, \xi_D)$ of two independent variables $\xi_{C,D} = (\boldsymbol{\rho}_{C,D} \cdot \mathbf{n})$, which in Gaussian case^{5,7} or in sharply peak approximation⁶ always contains a function $\Phi(\mp k\mathbf{n}) \propto \mathcal{H}(\zeta)$ of the one dimensionless variable $\zeta = (\mathbf{n}\mathbf{B}\mathbf{n}) > 0$ with the dimensionless symmetrical positively defined tensor \mathbf{B} built on the components of these vectors, whose traceless part \mathbf{B}_0 only is in fact necessary. Some technique with these functions is collected in Appendix B.

When $\Upsilon_1(k, \pm \mathbf{n}) < 0$, the last multiplier in expression (47) leads to suppression factor to the usual inverse square law for the event rate that reads as:

$$\propto \frac{1}{R^2} \left\{ 1 - \frac{\varrho_0^2}{R^2} \right\}, \quad \text{where: } \varrho_0^2 = -\frac{\overline{\Upsilon}_1(k, \pm \mathbf{n})}{(2k)^2}, \quad (48)$$

and the averaging $\overline{\Upsilon}_1$ relative to directions of $\boldsymbol{\rho}$ is implied for the case (a), that in fact approximately puts $\boldsymbol{\rho} = \mathbf{n}$, $\xi = 1$. For the case (b) one has to average relative to the vectors $\boldsymbol{\rho}_{C,D}$ independently, what makes the final result more model-dependent. According to Refs. 2,3 this suppression factor provided by the higher order corrections to Grimus-Stockinger formula (1) naturally explains the observed⁹ (anti-) neutrino deficit in reactor and others short-baseline experiments (see Refs. 2,3,9 and the references therein). Such explanation does not require any “new physics” and perhaps² can be tested experimentally.

Eqs. (32), (44) show that negative sign of coefficient Υ_1 originates by the most rapid variation $\Phi(\mp k\mathbf{n})$ with vector \mathbf{n} . Exponential variations are directly generated by the above limiting properties of the overlap function and are naturally arising in popular models of wave packets.⁵⁻⁷ Since beyond oscillation problem it is not meaningless to leave overlap function be defined by one wave packet only, any model

of wave packet also has to reproduce the above limiting properties of $\mathcal{F}_{C,D}(q)$.

In the spirit of Ref. 6 the following form of wave packet in momentum representation was suggested¹⁰ for $q_\mu \zeta_a^\mu \equiv (q\zeta_a) > 0$, with time-like 4-vector ζ_a^μ of mean quantum numbers of wave packet, such that $\zeta_a^0 > 0$, $\zeta_{a\mu} \zeta_a^\mu \equiv \zeta_a^2 > 0$:

$$\zeta_a(p_a, \sigma_a) = p_a g_1(m_a, \sigma_a) + s_a g_2(m_a, \sigma_a), \quad \phi^\sigma(\mathbf{q}, \mathbf{p}_a) = N_\sigma(m_a, \zeta_a^2) e^{-(q\zeta_a)}, \quad (49)$$

where both the momentum 4-vectors $q^\mu = (E_q, \mathbf{q})$, $p_a^\mu = (E_{p_a}, \mathbf{p}_a)$ are on mass shell: $E_q = \sqrt{\mathbf{q}^2 + m_a^2} > 0$, and so on, s_a - is a spin 4-vector, $g_1 \gg |g_2|$ - are some real functions¹⁰ of mass m_a and width σ_a , and $\aleph(\tau) m_a^{-2} = N_\sigma > 0$, is normalization constant with fixed asymptotic behavior¹⁰ as a function of dimensionless invariant variable $\tau = m_a \sqrt{\zeta_a^2(p_a, \sigma_a)}$, at $\tau \rightarrow \infty$ ($\sigma_a \rightarrow 0$), and at $\tau \rightarrow 0$ ($\sigma_a \rightarrow \infty$):

$$\aleph(\tau) \xrightarrow{\tau \rightarrow \infty} 2(2\pi)^{3/2} \tau^{3/2} e^\tau, \quad \text{and} \quad \aleph(\tau) \xrightarrow{\tau \rightarrow 0} \aleph(0) > 0. \quad (50)$$

Coordinate representation of this wave packet with fixed center x_a for scalar case satisfies Klein-Gordon equation and for $x^\mu = (t, \mathbf{x})$ is defined¹⁰ by Wightman function analytically continued¹⁵ into the same future tube (V^+ : $\zeta_a^0 > 0$, $\zeta_a^2 > 0$) as:

$$\begin{aligned} F_{p_a x_a}(x) &= e^{-i(p_a x_a)} \int \frac{d^3 \mathbf{q}}{(2\pi)^3 2E_q} \phi^\sigma(\mathbf{q}, \mathbf{p}_a) e^{-i(q(x-x_a))} = \\ &= (-i) e^{-i(p_a x_a)} N_\sigma D_{m_a}^-(x - x_a - i\zeta_a(p_a, \sigma_a)), \quad (g_2 \equiv 0). \end{aligned} \quad (51)$$

This wave packet conforms with general requirements of quantum field theory¹⁵ and due to (50) admits adequate description¹⁰ of both the limits to the states localized in momentum and coordinate spaces at $\sigma_a \rightarrow 0$ and $\sigma_a \rightarrow \infty$ respectively:

$$\phi^\sigma(\mathbf{q}, \mathbf{p}_a) \xrightarrow{\sigma \rightarrow 0} (2\pi)^3 2E_q \delta_3(\mathbf{q} - \mathbf{p}_a), \quad \phi^\sigma(\mathbf{q}, \mathbf{p}_a) \xrightarrow{\sigma \rightarrow \infty} N_\infty. \quad (52)$$

Furthermore, because $2(qp_a) = 2m_a^2 - (q - p_a)^2$, the nonrelativistic limit of (49), (51) for $g_1 = \sigma_a^{-2}$ exactly reproduces¹⁰ the usual Gaussian profiles in coordinate and momentum representations independently of normalization arbitrariness (50).

To get most simple example preserving oscillations we take an overlap function with one wave packet (49), (51) in the one vertex representing the another vertex by infinitely heavy nucleus in the spirit of Kobzarev model.¹⁶ For antineutrino this may be achieved for example in the Grimus-Stockinger model⁴ with nucleons in creation vertex $\{C\}$ fixed in infinitely heavy nucleus and the bound state of initial electron in detection vertex $\{D\}$ replaced by above wave packet state for free electron. For the case of neutrino one can take the decay $\pi^+ \rightarrow \mu^+ + \nu_\mu$ with wave packet (49) for pion state and plane wave for muon in $\{C\}$ - vertex taking $\{D\}$ - vertex as in the Kobzarev model.^{5,16} The overlap functions⁵ $\mathcal{F}_{C,D}(-p)$ for the first case (compare with Eq. (11) from Ref. 4 for $\tilde{J}_\lambda = \text{const}$), with $E_w = \sqrt{\mathbf{w}^2 + m_e^2}$, are reduced to:

$$\mathcal{F}_C(-p) = \text{const} \delta(p^0 - Q_C^0), \quad Q_C^0 = \Delta M - U^0, \quad Q_D = K - W, \quad (53)$$

$$\mathcal{F}_D(-p) = (2\pi)^4 \delta(p^0 + E_w - K^0) \frac{\phi_e^\sigma(\mathbf{w}, \mathbf{W})}{2E_w}, \quad \mathbf{w} = \mathbf{K} - \mathbf{p}, \quad W^0 = E_w, \quad (54)$$

where: ΔM is neutron-proton mass difference, $U^\mu = (U^0, \mathbf{U})$ is 4-momentum of created electron in vertex $\{C\}$; $W^\mu = (W^0, \mathbf{W})$ is the most probable 4-momentum of incoming initial electron with $\zeta_e(m_e, \sigma_e) = g_1 W$ in Eq. (49), $w^\mu = (E_w, \mathbf{w})$, and $K^\mu = (K^0, \mathbf{K})$ is the total outgoing 4-momentum carried away by final electron with final antineutrino in vertex $\{D\}$. Note that 3-vectors \mathbf{U} and \mathbf{Q}_C become indefinite for this case. The product of these functions belongs to $S(\mathbf{R}_p^3)$ only for replaced argument of second delta-function with $E_w \mapsto E_W$, which appears naturally in sharply peak approximation for $|\mathbf{p} - \mathbf{Q}_D| = |\mathbf{W} - \mathbf{w}| \ll m_e$, leading to:

$$\mathcal{F}(-p) \approx \text{const } \delta(p^0 - Q_C^0) \delta(p^0 - Q_D^0) \frac{N_{\sigma_e}}{2W^0} e^{-g_1(wW)}, \quad \text{where: } g_1 = \frac{1}{\sigma_e^2}, \quad (55)$$

$$(wW) = E_w W^0 - (\mathbf{w} \cdot \mathbf{W}) \approx m_e^2 + \frac{1}{2} ((\mathbf{p} - \mathbf{Q}_D)^j (\delta^{jl} - V^j V^l) (\mathbf{p} - \mathbf{Q}_D)^l) = \quad (56)$$

$$= m_e^2 + \frac{1}{2} \{ \mathbf{Q}_D^2 - (\mathbf{Q}_D \cdot \mathbf{V})^2 + \mathbf{p}^2 - 2(\mathbf{p} \cdot [\mathbf{Q}_D - \mathbf{V}(\mathbf{Q}_D \cdot \mathbf{V})]) - \mathbf{V}^2(\mathbf{p} \cdot \mathbf{p}) \}, \quad (57)$$

$$\text{for: } \mathbf{V} = \frac{\mathbf{W}}{W^0} = |\mathbf{V}| \boldsymbol{\omega}, \quad \omega^2 = 1, \quad B^{jl} = \omega^j \omega^l, \quad B_0^{jl} = \omega^j \omega^l - \frac{\delta^{jl}}{3}, \quad (58)$$

with vector \mathbf{V} as a most probable velocity of initial electron in detector. To define the coefficients $\Upsilon_s(k, \mathbf{n})$ (44) of asymptotic expansion (42) by making use of Eqs. (31)–(33), only the last two summands in the line (57) are now necessary, giving exactly the discussed here structure of the function $\Phi(\mathbf{q})$ (where due to the absence of vector \mathbf{Q}_C its role partially plays vector \mathbf{V}). For non-relativistic initial electron $|\mathbf{V}| \ll 1$, and neglecting this velocity, for $\mathbf{Q}_D = |\mathbf{Q}_D| \boldsymbol{\rho}$ and real $\lambda = g_1 k |\mathbf{Q}_D| > 0$, now it is enough to take (see Appendix B for details and another cases):

$$\Phi(\mathbf{q}) = \Phi(-\mathbf{p}) = \Phi(-k\mathbf{n}) \xrightarrow{|\mathbf{V}| \ll 1} e^{\lambda \xi}, \quad \xi = (\boldsymbol{\rho} \cdot \mathbf{n}) = \cos \Theta, \quad \text{whence:} \quad (59)$$

$$C_1(k, \mathbf{n}) = 2\lambda \xi - \lambda^2(1 - \xi^2), \quad \Upsilon_1(k, \mathbf{n}) = -4\lambda^2 [1 - (2 + \lambda \xi)(1 - \xi^2)]. \quad (60)$$

So $\Upsilon_1 < 0$, if $\lambda \xi < [(1 - \xi^2)^{-1} - 2]$, that $\forall \lambda > 0$ includes interval $3\pi/4 \leq \Theta \leq \pi$,

$$\text{which for } \lambda \rightarrow \infty, \text{ dilates to: } 0 \leq \Theta \leq 1/\sqrt{\lambda}, \quad \pi/2 + 1/\lambda \leq \Theta \leq \pi. \quad (61)$$

With the same function (59) $\forall \lambda < 0$ this condition implies interval $0 \leq \Theta \leq \pi/4$,

$$\text{dilating to: } 0 \leq \Theta \leq \pi/2 - 1/|\lambda|, \quad \pi - 1/\sqrt{|\lambda|} \leq \Theta \leq \pi, \quad \text{for } |\lambda| \rightarrow \infty, \quad (62)$$

that means the formal interchange in (61) of forward and backward hemispheres $\Theta \mapsto \pi - \Theta$. Similarly for the above mentioned case with neutrino, one can obtain for $\Phi(\mathbf{q}) = \Phi(k\mathbf{n})$ the same result (59) with $\lambda > 0$. Therefore for the both cases $\Upsilon_1(k, \mathbf{n}) < 0$ near the forward direction (61) with the narrow wave packet $\sigma_{e,\pi} \rightarrow 0$ and/or with the high energy $p^0 = Q_D^0$ of (anti-) neutrino. Since due to Eq. (56), only this region in fact contributes to the averaging relative to $\boldsymbol{\rho}$, from (60) for the parameter ϱ_0 (48), with $k = \sqrt{(Q_D^0)^2 - m_j^2} \approx |\mathbf{Q}_D|$, and $Q_D^0 = Q_C^0$ (53), one finds:

$$\overline{\Upsilon}_1 \simeq -4\lambda^2, \quad \varrho_0^2 \simeq \frac{4\lambda^2}{(2k)^2} = \frac{\mathbf{Q}_D^2}{\sigma_e^4}, \quad \varrho_0 \approx \frac{k}{\sigma_e^2}, \quad (63)$$

$$\text{where: } m_j \leq \sigma_e \ll m_e, \quad k \leq Q_C^0 \leq \Delta M - m_e. \quad (64)$$

For $k \simeq \Delta M - m_e \approx 0.783 \text{ MeV}$ and $\sigma_e \sim m_j \simeq 0.2 \text{ eV}$ it follows $\varrho_0 \simeq 3.86 \text{ m}$. Note that for real β -decay of heavy nucleus the value of antineutrino energy $Q_C^0 \simeq k$ may be in many times greater. This qualitative analytical estimation illustrates and elucidates the numerical calculations^{2,3} with more detailed models, that give the best value for this parameter $\varrho_0 \simeq 3.5 \text{ m}$. The obtained analytical results (31)–(33), (42), (44), together with relations from Appendix B can essentially simplify such calculations making them probably more transparent and model-independent.

5. Conclusions

As shown in Appendix A, following the same way as in Lemma 1 the operator expansion (20) for the Green function of Helmholtz equation may be generalized for arbitrary N -dimensional Euclidean space. The same concerns the assertion of Theorem 1 about asymptotic expansion for N -dimensional analog of integral (1).

The following comments are in order. Since the main principles of quantum field theory¹⁵ require the wave packets belong to $S(\mathbf{R}_q^3)$, the same holds true for their convolution and thus for $\Phi(\mathbf{q})$, as was supposed in Theorem 1. This property ensures the possibility of further asymptotic expansion (31), (37) of integral (1) and defines the order of leading corrections to Grimus-Stockinger asymptotic,⁴ which is crucial for explanation^{2,3} (48) of (anti-) neutrino deficit.

For electron in stationary bound state⁴ with energy $W^0 = E_b$ instead of the wave packet state the overlap function (54), with the also indefinite now vectors \mathbf{W} , \mathbf{K} and \mathbf{Q}_D , becomes: $\mathcal{F}_D(-p) = \delta(p^0 + E_b - K^0) \tilde{\psi}_b(\mathbf{w}) \in C^\infty(\mathbf{R}_p^3)$, where $|\tilde{\psi}_b(\mathbf{K} - \mathbf{p})| \leq O(|\mathbf{p}|^{-4})$ for $|\mathbf{p}| \rightarrow \infty$. It may be argued following to the Lemma 1 of Ref. 2 and the Theorem of Ref. 4, that this fourth power fall-off should be enough for the well definiteness of the asymptotic expansion (31) up to the term with $s = 2$, and therefore the expansion (42), (44) up to the term with $n = 1$, what preserves Eqs. (47), (48). On the other hand the infinite coherence length of neutrino oscillations for the stationary case⁴ is also the price for the absence of external wave packets like (49), (51); only their presence in both the creation $\{C\}$ and detection $\{D\}$ vertices provides a finite value of coherence length.^{5,6} So we conclude, that again as in the Ref. 4: “Therefore the physics under consideration seems to comply naturally with the mathematical requirements”.

Since the wave packet (49) in momentum representation $\phi^\sigma(\mathbf{q}, \mathbf{p}_a) \in S(\mathbf{R}_q^3)$, the same takes place¹⁵ for its coordinate representation (51). So, for fixed t , $F_{p_a x_a}(t, \mathbf{x}) \in S(\mathbf{R}_x^3)$ obviously with infinite support and the same holds true for their convolution $\phi(\mathbf{x})$ in Eqs. (36) and (12). Therefore the expansion (37) acquires also the additional asymptotic nature, with nonzero corrections (38), governed by detailed form of asymptotic behavior of the function $\phi(\mathbf{x})$ itself. These corrections inevitable introduce an additional dimensional parameter and may also affect the behavior of $\mathcal{J}(\mathbf{R})$ at microscopically big but macroscopically small distances R , discussed here and in Ref. 2,3,9.

Appendix A.

For N -dimension space, defining: $\nu = (N - 2)/2$, $a = (N - 3)/2 = \nu - \frac{1}{2}$, $l = 0, 1, 2, \dots$, $j = l + a$, with: $l(l + 2\nu) + \nu^2 - \frac{1}{4} = l(l + 2\nu) + a(a + 1) = j(j + 1)$, for $\mathbf{R} = R\mathbf{n}$, $\mathbf{x} = r\mathbf{v}$, $\mathbf{k} = k\mathbf{n}$, one has^{12, 13} Gegenbauer polynomials $C_l^{(\nu)}((\mathbf{n} \cdot \mathbf{v}))$ as eigenfunction of N -dimension square angular momenta operator $\mathcal{L}_{\mathbf{n}}^{(N)}$, where for $\nabla_{\mathbf{R}}$ in (4), instead of (5), (6): $[(\partial_{\mathbf{n}})_{\beta}, n_{\alpha}] = \delta_{\alpha\beta} - n_{\alpha}n_{\beta}$, $(\mathbf{n} \cdot \partial_{\mathbf{n}}) = 0$, $(\partial_{\mathbf{n}} \cdot \mathbf{n}) = N - 1$, $x_{\alpha}\partial_{\beta} - x_{\beta}\partial_{\alpha} = n_{\alpha}(\partial_{\mathbf{n}})_{\beta} - n_{\beta}(\partial_{\mathbf{n}})_{\alpha} = iL_{\alpha\beta}$ and $-\partial_{\mathbf{n}}^2 = \mathbf{L}^2/2 = \mathcal{L}_{\mathbf{n}}^{(N)}$, so that:¹³

$$\nabla_{\mathbf{R}}^2 = \partial_R^2 + \frac{N-1}{R}\partial_R - \frac{\mathcal{L}_{\mathbf{n}}^{(N)}}{R^2} = \frac{1}{R^{a+1}}\partial_R^2 R^{a+1} - \frac{a(a+1)}{R^2} - \frac{\mathcal{L}_{\mathbf{n}}^{(N)}}{R^2}, \quad (\text{A.1})$$

$$\mathcal{L}_{\mathbf{n}}^{(N)} C_l^{(\nu)}((\mathbf{n} \cdot \mathbf{v})) = l(l + 2\nu) C_l^{(\nu)}((\mathbf{n} \cdot \mathbf{v})). \quad (\text{A.2})$$

In definitions (27), (28), where,^{8, 11–13} for $l \mapsto j$, with $|\arg z - \beta_{1,2}| < \pi/2$:

$$K_{\lambda}(z) = \frac{1}{2} \int_{0e^{i\beta_2}}^{\infty e^{-i\beta_1}} \frac{dt}{t} t^{\pm\lambda} \exp\left\{-\frac{z}{2}\left(t + \frac{1}{t}\right)\right\}, \quad (\text{A.3})$$

$$\text{and: } \int_0^{\infty} dr \psi_{j0}(kr) \psi_{j0}(qr) = \frac{\pi}{2} \delta(q - k), \quad (\text{A.4})$$

according to formulas (8.534), (8.532) of Ref. 12 ($H_{\nu}^{(1),(2)}(z)$ are the Hankel functions,^{8, 12}) the plane wave and the Green function now read:^{13, 17}

$$e^{\mp i(\mathbf{k} \cdot \mathbf{x})} = \sqrt{\frac{2}{\pi}} \frac{2^{\nu-1} \Gamma(\nu)}{(\mp ikr)^a k r} \sum_{j=a}^{\infty} (2j+1) i^{\mp j} \psi_{j0}(kr) C_l^{(\nu)}((\mathbf{n} \cdot \mathbf{v})), \quad (\text{A.5})$$

$$\left\langle \mathbf{R} \left| \frac{1}{-\nabla^2 - k^2 \mp i0} \right| \mathbf{x} \right\rangle = \int \frac{d^N \mathbf{q}}{(2\pi)^N} \frac{e^{i(\mathbf{q} \cdot (\mathbf{R} - \mathbf{x}))}}{(\mathbf{q}^2 - k^2 \mp i0)} = \quad (\text{A.6})$$

$$= \pm \frac{i}{4} \left(\frac{k}{2\pi}\right)^{\nu} \frac{H_{\nu}^{(2)}(k|\mathbf{R} - \mathbf{x}|)}{|\mathbf{R} - \mathbf{x}|^{\nu}} = \sqrt{\frac{\pi}{2}} \frac{(\mp i k)^a}{(2\pi)^{\nu+1}} \frac{\chi_a(\mp i k |\mathbf{R} - \mathbf{x}|)}{|\mathbf{R} - \mathbf{x}|^{a+1}} = \quad (\text{A.7})$$

$$= \frac{\Gamma(\nu)}{4\pi^{\nu+1}} \frac{1}{(Rr)^{a+1}} \sum_{j=a}^{\infty} (2j+1) \frac{i^{\mp j}}{k} \chi_j(\mp i k R) \psi_{j0}(kr) C_l^{(\nu)}((\mathbf{n} \cdot \mathbf{v})), \quad (\text{A.8})$$

for $R > r$ and $l = j - a$, where a and $j \geq a$ are both integer for odd N or half integer for even N . So, for $R > r$, with again really not appearing operator $l \mapsto \Lambda_{\mathbf{n}}^{(N)}$, one obtains similarly the following generalization of Lemma 1 (19), (20):

$$\sqrt{\frac{\pi}{2}} \frac{(\mp i k)^a}{(2\pi)^{\nu+1}} \frac{\chi_a(\mp i k |\mathbf{R} - \mathbf{x}|)}{|\mathbf{R} - \mathbf{x}|^{a+1}} = \sqrt{\frac{\pi}{2}} \frac{(\mp i k)^a}{(2\pi)^{\nu+1}} \frac{\chi_{\Lambda_{\mathbf{n}}^{(N)} + a}(\mp i k R)}{R^{a+1}} e^{\mp i k (\mathbf{n} \cdot \mathbf{x})} \sim \quad (\text{A.9})$$

$$\sim \sqrt{\frac{\pi}{2}} \frac{(\mp i k)^a}{(2\pi)^{\nu+1}} \frac{e^{\pm i k R}}{R^{a+1}} \left\{ 1 + \sum_{s=1}^{\infty} \frac{\prod_{\mu=1}^s [\mathcal{L}_{\mathbf{n}}^{(N)} + a(a+1) - \mu(\mu-1)]}{s! (\mp 2 i k R)^s} \right\} e^{\mp i k (\mathbf{n} \cdot \mathbf{x})}. \quad (\text{A.10})$$

Here (A.9) is again an operator rewrite of (A.8) via (A.5) and positively defined self-adjoint operator with positive eigenvalue, from which $\chi_j(z)$ (27) depends on again as a whole function (A.3):

$$\mathcal{L}_{\mathbf{n}}^{(N)} + a(a+1) + \frac{1}{4} = \left(\Lambda_{\mathbf{n}}^{(N)} + a + \frac{1}{2} \right)^2 \mapsto \left(j + \frac{1}{2} \right)^2, \quad (\text{A.11})$$

and again the asymptotic version^{8,12} of expansion (27) is used. Due to (A.1), instead of operator $\mathcal{L}_{\mathbf{n}}^{(N)} + a(a+1)$ in (A.10), again may be used the operator from the square brackets of the radial Schrödinger equation:

$$\left[r^2 \left(\frac{1}{r^{a+1}} \partial_r^2 r^{a+1} + k^2 \right) \right] \frac{\psi_{j0}(kr)}{r^{a+1}} = j(j+1) \frac{\psi_{j0}(kr)}{r^{a+1}}. \quad (\text{A.12})$$

Appendix B.

The differential operators $\partial_{\mathbf{n}}$ and $\mathcal{L}_{\mathbf{n}} = -\partial_{\mathbf{n}}^2$ defined in Eqs. (4)–(7) have the following properties for any differentiable scalar functions $f(\xi)$, $\mathcal{W}(\xi, \varsigma)$, $\mathcal{H}(\zeta)$ with ξ from Eq. (59), $\varsigma = (\boldsymbol{\rho} \cdot \mathbf{n})$, $\zeta_m = (\mathbf{n} \mathbf{B}^m \mathbf{n})$, $\zeta \equiv \zeta_1$, $Tr\{\mathbf{B}^m\} = 3\bar{\zeta}_m$, $\zeta_{m0} = \zeta_m - \bar{\zeta}_m$:

$$\partial_{\mathbf{n}} f(\xi) = \partial_{\mathbf{n}} f((\boldsymbol{\rho} \cdot \mathbf{n})) = \boldsymbol{\rho}_{\perp} \partial_{\xi} f(\xi) + f(\xi) \partial_{\mathbf{n}}, \quad \boldsymbol{\rho}_{\perp} = \boldsymbol{\rho} - \mathbf{n}(\boldsymbol{\rho} \cdot \mathbf{n}), \quad (\text{B.1})$$

$$(\partial_{\mathbf{n}} \xi) = \boldsymbol{\rho}_{\perp}, \quad (\partial_{\mathbf{n}} \varsigma) = \boldsymbol{\rho}_{\perp}, \quad ((\partial_{\mathbf{n}})^j \zeta_m) = 2 [(\mathbf{B}^m \mathbf{n})^j - \mathbf{n}^j \zeta_m], \quad (\text{B.2})$$

$$\mathcal{L}_{\mathbf{n}} \xi = 2\xi, \quad \mathcal{L}_{\mathbf{n}} \varsigma = 2\varsigma, \quad \mathcal{L}_{\mathbf{n}} \zeta_m = 2[3\zeta_m - Tr\{\mathbf{B}^m\}] \equiv 6\zeta_{m0} = \mathcal{L}_{\mathbf{n}} \zeta_{m0}, \quad (\text{B.3})$$

$$f(\xi) C_1(\xi) = \mathcal{L}_{\mathbf{n}} f(\xi) = \partial_{\xi}(\xi^2 - 1) \partial_{\xi} f(\xi) = [(\xi^2 - 1) \partial_{\xi}^2 + 2\xi \partial_{\xi}] f(\xi), \quad (\text{B.4})$$

$$f(\xi) C_2(\xi) = \left[(6\xi^2 - 2) \partial_{\xi}^2 + 4\xi(\xi^2 - 1) \partial_{\xi}^3 + \frac{1}{2}(\xi^2 - 1)^2 \partial_{\xi}^4 \right] f(\xi), \quad (\text{B.5})$$

$$\mathcal{L}_{\mathbf{n}} \mathcal{W}(\xi, \varsigma) = [\partial_{\xi}(\xi^2 - 1) \partial_{\xi} + \partial_{\varsigma}(\varsigma^2 - 1) \partial_{\varsigma} + 2(\xi \varsigma - (\boldsymbol{\rho} \cdot \boldsymbol{\rho})) \partial_{\xi \varsigma}^2] \mathcal{W}(\xi, \varsigma), \quad (\text{B.6})$$

$$\mathcal{L}_{\mathbf{n}} f(\xi) g(\varsigma) = g(\varsigma) \mathcal{L}_{\mathbf{n}} f(\xi) + f(\xi) \mathcal{L}_{\mathbf{n}} g(\varsigma) - 2(\boldsymbol{\rho}_{\perp} \cdot \boldsymbol{\rho}_{\perp}) \partial_{\xi} f(\xi) \partial_{\varsigma} g(\varsigma), \quad (\text{B.7})$$

$$\mathcal{H}(\zeta) C_1(\zeta) = \mathcal{L}_{\mathbf{n}} \mathcal{H}(\zeta) = \{6\zeta_0 \partial_{\zeta} + 4(\zeta^2 - \zeta_2) \partial_{\zeta}^2\} \mathcal{H}(\zeta), \quad \zeta_0 \equiv \zeta_{10}, \quad (\text{B.8})$$

$$\mathcal{H}(\zeta) C_2(\zeta) = \frac{1}{2} [\mathcal{L}_{\mathbf{n}}^2 - 2\mathcal{L}_{\mathbf{n}}] \mathcal{H}(\zeta) = \quad (\text{B.9})$$

$$= \{12\zeta_0 \partial_{\zeta} + 6[(4\zeta + 3\zeta_0)\zeta_0 + 6(\zeta^2 - \zeta_2) - 2\zeta_{20}] \partial_{\zeta}^2 + 8[(4\zeta + 3\zeta_0)(\zeta^2 - \zeta_2) + 2(\zeta_3 - \zeta\zeta_2)] \partial_{\zeta}^3 + 8(\zeta^2 - \zeta_2)^2 \partial_{\zeta}^4\} \mathcal{H}(\zeta). \quad (\text{B.10})$$

When $f(\xi) \mapsto e^{\lambda \xi}$, Eqs. (B.4), (B.5) with $\partial_{\xi} \mapsto \lambda$ immediately give Eqs. (60). Similarly for $\mathcal{H}(\zeta) \mapsto e^{\lambda \zeta}$ one has $\partial_{\zeta} \mapsto \lambda$ in Eqs. (B.8)–(B.10), leading to:

$$\Upsilon_1 = -8\lambda [a\lambda^2 + b\lambda + c], \quad \text{with: } a = 4[2\zeta^3 + \zeta_3 - 3\zeta\zeta_2], \quad (\text{B.11})$$

$$b = 3[2\zeta_0\zeta + 3(\zeta^2 - \zeta_2) - \zeta_{20}], \quad c = 3\zeta_0, \quad \zeta_m = (\mathbf{n} \mathbf{B}^m \mathbf{n}) > 0, \quad (\text{B.12})$$

$$(\zeta_m)^2 \leq \zeta_{2\sigma} \zeta_{2m-2\sigma}, \quad \zeta_m \geq \zeta_{m-1} \geq \zeta^2 \zeta_{m-2} \geq \dots \geq \zeta^k \zeta_{m-k} \dots \geq \zeta^m, \quad (\text{B.13})$$

due to positive definiteness of \mathbf{B} and Cauchy-Schwarz inequality. Formally the case (B.11), (B.12) appears in Eqs. (57) for “collinear” events with $\mathbf{K} \parallel \mathbf{V}$ only, for $\lambda = g_1 k^2 \mathbf{V}^2 / 2 > 0$ and degenerate tensor \mathbf{B} (58) evidently realizing all inequalities (B.13). The structure of \mathbf{B} and λ becomes much more complicated for the complete

models.^{2,6} Nevertheless the most general case containing all variables simultaneously, like $\Phi(\mp k\mathbf{n}) \mapsto e^{\alpha\xi}e^{\beta\varsigma}e^{\lambda\zeta}$, may be considered by the same way using the above relations together.

Acknowledgments

Authors thank V. Naumov, D. Naumov, and N. Ilyin for useful discussions and Reviewer for constructive comments.

References

1. S. E. Korenblit, D. V. Taychenachev, *Phys. Part. Nuc. Let.* **10** 1002 (2013). (arXiv: 1304.5192 [hep-th])
2. V. A. Naumov, D. S. Shkirmanov, *Eur. Phys. J. C* **73** 2627 (2013). (arXiv: 1309.1011 [hep-th])
3. D. V. Naumov, V. A. Naumov, D. S. Shkirmanov, *Neutrino flavor transitions in quantum field theory and generalized Grimus-Stockinger theorem*, in *Rep. on 41-st ITEP Int. Winter School of Phys.*, (Moscow, Russia, 2013). (http://www.itep.ru/ws/2013/lec/YSS/Feb15/Feb15/5-prezentaciya_itep.pdf)
4. W. Grimus, P. Stockinger, *Phys. Rev. D* **54** 3414 (1996).
5. M. Beuthe, *Phys. Rep.* **375** 105 (2003).
6. V. A. Naumov, D. V. Naumov, *J. Phys. G: Nucl. Part. Phys.* **37** 105014 (2010). (arXiv: 1008.0306v2 [hep-ph])
7. E. Kh. Akhmedov, J. Kopp, *JHEP* **04** 008 (2010).
8. F. Olver, *Introduction to asymptotics and special function*. (Acad. Press, NY, 1974).
9. G. Mention, M. Fechner, Th. Lasserre, Th.A. Mueller, D. Lhuillier, et al., *Phys. Rev. D* **83** 073006 (2011). (arXiv: 1101.2755 [hep-ex])
10. S. E. Korenblit, D. V. Taychenachev, *On uniqueness of the wave packet in QFT and neutrino oscillation problem*, in *Proc. of XIII Baikal School on Fund. Phys.: Interaction of Fields and Radiation with Matter*, (eds. E. V. Devyatova, ISEP, Irkutsk, Russia, 2013). pp 109-114. (in Russian) (<http://bsfp.iszf.irk.ru/images/2013/Sbornik-2013/Section-A/Taychenachev-109-114.pdf>)
11. J. R. Taylor, *Scattering Theory*. (J. Wiley & Sons Inc. NY, 1972).
12. I. S. Gradshteyn, I. M. Rizhik, *Tables of Integrals, Sums, Series and Products*, 7th edition, (Acad. Press, San Diego, 2007).
13. N. Ya. Vilenkin, *Special Functions and Group Representations Theory*. (Nauka, Moscow, 1991). (in Russian)
14. L. C. Biedenharn, J. D. Louck, *Angular Momentum in Quantum Physics. Theory and Application*. (Addison-Wesley Pub. Com., Reading, Massachusetts, 1981).
15. N. N. Bogoliubov, A. A. Logunov, I. T. Todorov, *Foundation of Axiomatic Approach in Quantum Field Theory*. (Nauka, Moscow, 1969). (in Russian).
16. I. Yu. Kobzarev, B. V. Martemyanov, L.B.Okun, M.G.Shchepkin, *Sov. J. Nucl. Phys.* **35** 708 (1982) [*Yad. Fiz.* **35** 1210 (1982)].
17. S. E. Korenblit, Yu. V. Parfenov, *Phys. Atom. Nucl.*, **56** 483 (1993), [*Yad. Fiz.* **56** 105 (1993)], (arXiv: 0901.0341 [hep-th])